



YEAR 2008

[Insulin sensitivity and resistin levels in gestational diabetes mellitus and after parturition.](#) Megia A, Vendrell J et al. Eur J Endocrinol. 2008 Feb;158(2):173-8. doi: 10.1530/EJE-07-0671.

[Adult subjects with Prader-Willi syndrome show more low-grade systemic inflammation than matched obese subjects.](#) Caixàs A, Giménez-Palop O et al. J Endocrinol Invest. 2008 Feb;31(2):169-75.

[Human serum levels of fetal antigen 1 \(FA1/Dlk1\) increase with obesity, are negatively associated with insulin sensitivity and modulate inflammation in vitro.](#) Chacón MR, Miranda M et al. Epub 2008 Apr 8.

[Influence of morbid obesity and insulin resistance on gene expression levels of AQP7 in visceral adipose tissue and AQP9 in liver.](#) Catalán V, Gómez-Ambrosi J et al. Obes Surg. 2008 Jun;18(6):695-701. doi: 10.1007/s11695-008-9453-7.

[The IL-6 system in HIV-1-infection and in HAART-related fat redistribution syndromes.](#) Saumoy M, López-Dupla M et al. AIDS. 2008 Apr 23;22(7):893-6. doi: 10.1097/QAD.0b013e3282f4dde7.

[Regulation of bone mineral density in morbidly obese women: a cross-sectional study in two cohorts before and after bypass surgery.](#) Gómez JM, Vilarrasa N et al. Obes Surg. 2009 Mar;19(3):345-50. doi: 10.1007/s11695-008-9529-4.

[The nuclear receptor coactivator AIB3 is a modulator of HOMA beta-cell function in nondiabetic children.](#) Burghardt H, López-Bermejo A et al. Clin Endocrinol (Oxf). 2008 Nov;69(5):730-6. doi: 10.1111/j.1365-2265.2008.03232.x.

[LMNA mRNA expression is altered in human obesity and type 2 diabetes.](#) Miranda M, Chacón MR et al. Obesity (Silver Spring). 2008 Aug;16(8):1742-8. doi: 10.1038/oby.2008.276.

[Circulating soluble CD36 is a novel marker of liver injury in subjects with altered glucose tolerance.](#) Fernández-Real JM, Handberg A et al. J Nutr Biochem. 2009 Jun;20(6):477-84. doi: 10.1016/j.jnutbio.2008.05.009.

[Urinary interleukin-6 excretion reflects mean 24-hour systolic blood pressure in type 2 diabetes.](#) Esteve E, Torquet P et al. Endocrinol Nutr. 2008 Oct;55(9):383-8. doi: 10.1016/S1575-0922(08)75074-5.