



YEAR 2009

[Regulation of bone mineral density in morbidly obese women: a cross-sectional study in two cohorts before and after bypass surgery.](#) Gómez JM, Vilarrasa N et al. *Obes Surg.* 2009 Mar;19(3):345-50. doi: 10.1007/s11695-008-9529-4.

[Joint analysis of individual participants' data from 17 studies on the association of the IL6 variant -174G>C with circulating glucose levels, interleukin-6 levels, and body mass index.](#) Huth C, Illig T et al. *Ann Med.* 2009;41(2):128-38. doi: 10.1080/07853890802337037.

[Circulating soluble CD36 is a novel marker of liver injury in subjects with altered glucose tolerance.](#) Fernández-Real JM, Handberg A et al. *J Nutr Biochem.* 2009 Jun;20(6):477-84. doi: 10.1016/j.jnutbio.2008.05.009.

[Urinary interleukin-6 excretion reflects mean 24-hour systolic blood pressure in type 2 diabetes.](#) Esteve E, Torguet P et al. *Endocrinol Nutr.* 2008 Oct;55(9):383-8. doi: 10.1016/S1575-0922(08)75074-5.

[Circulating soluble CD36 is associated with glucose metabolism and interleukin-6 in glucose-intolerant men.](#) Handberg A, Lopez-Bermejo A et al. *Diab Vasc Dis Res.* 2009 Jan;6(1):15-20. doi: 10.3132/dvdr.2009.003.

[Adipocyte fatty acid-binding protein as a determinant of insulin sensitivity in morbid-obese women.](#) Simón I, Escoté X et al. *Obesity (Silver Spring).* 2009 Jun;17(6):1124-8. doi: 10.1038/oby.2008.665.

[Plasma visfatin concentrations increase in both hyper and hypothyroid subjects after normalization of thyroid function and are not related to insulin resistance, anthropometric or inflammatory parameters.](#) Caixàs A, Tirado R et al. *Clin Endocrinol (Oxf).* 2009 Nov;71(5):733-8. doi: 10.1111/j.1365-2265.2009.03546.x.

[PPARgamma Pro12Ala polymorphism in HIV-1-infected patients with HAART-related lipodystrophy.](#) Saumoy M, Veloso S et al. *Curr HIV Res.* 2009 Sep;7(5):533-40.

[Grape-seed procyanidins modulate inflammation on human differentiated adipocytes in vitro.](#) Chacón MR, Ceperuelo-Mallafré V et al. *Cytokine.* 2009 Aug;47(2):137-42. doi: 10.1016/j.cyto.2009.06.001.

[Gene expression of paired abdominal adipose AQP7 and liver AQP9 in patients with morbid obesity: relationship with glucose abnormalities.](#) Miranda M, Ceperuelo-Mallafré V et al. *Metabolism.* 2009 Dec;58(12):1762-8. doi: 10.1016/j.metabol.2009.06.004.

[Circulating and adipose tissue gene expression of zinc-alpha2-glycoprotein in obesity: its relationship with adipokine and lipolytic gene markers in subcutaneous and visceral fat.](#) Ceperuelo-Mallafré V, Näf S et al. *J Clin Endocrinol Metab.* 2009 Dec;94(12):5062-9. doi: 10.1210/jc.2009-0764.



DIABETES AND METABOLIC
ASSOCIATED DISEASES
RESEARCH GROUP

[Study of the relationship between adiponectin, interleukin-18, ghrelin and bone mineral density in morbidly obese women after gastric bypass.](#) Vilarrasa N, Gómez JM et al. Endocrinol Nutr. 2009 Aug-Sep;56(7):355-60. doi: 10.1016/S1575-0922(09)72454-4. Spanish.

[Metabolic endotoxemia and saturated fat contribute to circulating NGAL concentrations in subjects with insulin resistance.](#) Moreno-Navarrete JM, Manco M et al. Int J Obes (Lond). 2010 Feb;34(2):240-9. doi: 10.1038/ijo.2009.242.