



## YEAR 2015

[HIV-1/HAART-Related Lipodystrophy Syndrome \(HALS\) Is Associated with Decreased Circulating sTWEAK Levels.](#) López-Dupla M, Maymó-Masip E et al. PLoS One. 2015 Dec 14;10(12):e0144789. doi: 10.1371/journal.pone.0144789.

[Hypoxia is associated with a lower expression of genes involved in lipogenesis in visceral adipose tissue.](#) García-Fuentes E, Santiago-Fernández C et al. J Transl Med. 2015 Nov 30;13:373. doi: 10.1186/s12967-015-0732-5.

[Circulating Stem Cells Associate With Adiposity and Future Metabolic Deterioration in Healthy Subjects.](#) Fadini GP, Bonora BM et al. J Clin Endocrinol Metab. 2015 Dec;100(12):4570-8. doi: 10.1210/jc.2015-2867.

[FGF-23/Vitamin D Axis in Type 1 Diabetes: The Potential Role of Mineral Metabolism in Arterial Stiffness.](#) Llauradó G, Megia A et al. PLoS One. 2015 Oct 13;10(10):e0140222. doi: 10.1371/journal.pone.0140222.

[Adipose tissue glycogen accumulation is associated with obesity-linked inflammation in humans.](#) Ceperuelo-Mallafré V, Ejarque M et al. Mol Metab. 2015 Oct 16;5(1):5-18. doi: 10.1016/j.molmet.2015.10.001.

[Obesity changes the human gut mycobiome.](#) Rodríguez M, Pérez D et al. Sci Rep. 2015 Oct 12;5:14600. doi: 10.1038/srep14600.

[PPP2R5C Couples Hepatic Glucose and Lipid Homeostasis.](#) Cheng YS, Seibert O et al. PLoS Genet. 2015 Oct 6;11(10):e1005561. doi: 10.1371/journal.pgen.1005561.

[Differences in the osteogenic differentiation capacity of omental adipose derived stem cells in obese patients with and without metabolic syndrome.](#) Oliva-Olivera W, Gea AL et al. Endocrinology. 2015 Dec;156(12):4492-501. doi: 10.1210/en.2015-1413.

[Lipopolysaccharide binding protein is an adipokine involved in the resilience of the mouse adipocyte to inflammation.](#) Moreno-Navarrete JM, Escoté X et al. Diabetologia. 2015 Oct;58(10):2424-34. doi: 10.1007/s00125-015-3692-7.

[Haptoglobin genotype is associated with increased endothelial dysfunction serum markers in type 1 diabetes.](#) Llauradó G, Gutiérrez C et al. Eur J Clin Invest. 2015 Sep;45(9):932-9. doi: 10.1111/eci.12487.

[Zinc- \$\alpha\$ 2-Glycoprotein Modulates AKT-Dependent Insulin Signaling in Human Adipocytes by Activation of the PP2A Phosphatase.](#) Ceperuelo-Mallafré V, Ejarque M et al. PLoS One. 2015 Jun 11;10(6):e0129644. doi: 10.1371/journal.pone.0129644.

[HIV/antiretroviral therapy-related lipodystrophy syndrome \(HALS\) is associated with higher RBP4 and lower omentin in plasma.](#) Peraire J, López-Dupla M et al. Clin Microbiol Infect. 2015 Jul;21(7):711.e1-8. doi: 10.1016/j.cmi.2015.04.002.



DIABETES AND METABOLIC  
ASSOCIATED DISEASES  
RESEARCH GROUP

[Low blood levels of sTWEAK are related to locoregional failure in head and neck cancer.](#) Avilés-Jurado FX, Terra X et al. Eur Arch Otorhinolaryngol. 2015 Jul;272(7):1733-41. doi: 10.1007/s00405-014-3095-2.

[PPAR \$\beta/\delta\$  ameliorates fructose-induced insulin resistance in adipocytes by preventing Nrf2 activation.](#) Barroso E, Rodríguez-Rodríguez R et al. Biochim Biophys Acta. 2015 May;1852(5):1049-58. doi: 10.1016/j.bbadi.2015.02.010.

[Enhanced fatty acid oxidation in adipocytes and macrophages reduces lipid-induced triglyceride accumulation and inflammation.](#) Malandrino MI, Fuchio R et al. Am J Physiol Endocrinol Metab. 2015 May 1;308(9):E756-69. doi: 10.1152/ajpendo.00362.2014.

[Cord blood FGF21 in gestational diabetes and its relationship with postnatal growth.](#) Megia A, Gil-Lluis P et al. Acta Diabetol. 2015 Aug;52(4):693-700. doi: 10.1007/s00592-014-0705-9.

[Estimating Cardiovascular Risk in Spain by the European Guidelines on Cardiovascular Disease Prevention in Clinical Practice.](#) Amor AJ, Masana L et al. Rev Esp Cardiol (Engl Ed). 2015 May;68(5):417-25. doi: 10.1016/j.rec.2014.05.023.

[Reduced circulating levels of TWEAK are associated with gestational diabetes mellitus.](#) Simón-Muela I, Llauradó G et al. Eur J Clin Invest. 2015 Jan;45(1):27-35. doi: 10.1111/eci.12375.