



YEAR 2014

[Reduced circulating sTWEAK levels are associated with metabolic syndrome in elderly individuals at high cardiovascular risk.](#) Díaz-López A, Bulló M et al. *Cardiovasc Diabetol.* 2014 Feb 24;13:51. doi: 10.1186/1475-2840-13-51.

[Serum activin A and follistatin levels in gestational diabetes and the association of the Activin A-Follistatin system with anthropometric parameters in offspring.](#) Näf S, Escote X et al. *PLoS One.* 2014 Apr 24;9(4):e92175. doi: 10.1371/journal.pone.0092175.

[Disruption of GIP/GIPR axis in human adipose tissue is linked to obesity and insulin resistance.](#) Ceperuelo-Mallafré V, Duran X et al. *J Clin Endocrinol Metab.* 2014 May;99(5):E908-19. doi: 10.1210/jc.2013-3350.

[Prevalence of plasma lipid abnormalities and its association with glucose metabolism in Spain: the Di@bet.es study.](#) Martínez-Hervas S, Carmena R et al. *Clin Investig Arterioscler.* 2014 May-Jun;26(3):107-14. doi: 10.1016/j.arteri.2013.12.001.

[Serum sCD163 levels are associated with type 2 diabetes mellitus and are influenced by coffee and wine consumption: results of the Di@bet.es study.](#) Rojo-Martínez G, Maymó-Masip E et al. *PLoS One.* 2014 Jun 30;9(6):e101250. doi: 10.1371/journal.pone.0101250.

[Prevalence of obesity, diabetes and other cardiovascular risk factors in Andalusia \(southern Spain\). Comparison with national prevalence data. The Di@bet.es study.](#) Valdés S, García-Torres F et al. *Rev Esp Cardiol (Engl Ed).* 2014 Jun;67(6):442-8. doi: 10.1016/j.rec.2013.09.029.

[Advanced glycation end products are associated with arterial stiffness in type 1 diabetes.](#) Llauradó G, Ceperuelo-Mallafré V et al. *J Endocrinol.* 2014 Jun;221(3):405-13. doi: 10.1530/JOE-13-0407.

[Low blood levels of sTWEAK are related to locoregional failure in head and neck cancer.](#) Avilés-Jurado FX, Terra X et al. *Eur Arch Otorhinolaryngol.* 2015 Jul;272(7):1733-41. doi: 10.1007/s00405-014-3095-2.

[CCNG2 and CDK4 is associated with insulin resistance in adipose tissue.](#) Garrido-Sánchez L, Roca-Rodríguez Mdel M et al. *Surg Obes Relat Dis.* 2014 Jul-Aug;10(4):691-6. doi: 10.1016/j.soard.2013.12.011.

[Skeletal muscle myogenesis is regulated by G protein-coupled receptor kinase 2.](#) García-Guerra L, Vila-Bedmar R et al. *J Mol Cell Biol.* 2014 Aug;6(4):299-311. doi: 10.1093/jmcb/mju025.

[Variable patterns of obesity and cardiometabolic phenotypes and their association with lifestyle factors in the Di@bet.es study.](#) Gutiérrez-Repiso C, Soriguer F et al. *Nutr Metab Cardiovasc Dis.* 2014 Sep;24(9):947-55. doi: 10.1016/j.numecd.2014.04.019.



DIABETES AND METABOLIC
ASSOCIATED DISEASES
RESEARCH GROUP

[Human aquaporin-11 is a water and glycerol channel and localizes in the vicinity of lipid droplets in human adipocytes.](#) Madeira A, Fernández-Veledo S et al. *Obesity* (Silver Spring). 2014 Sep;22(9):2010-7. doi: 10.1002/oby.20792.

[Ambient temperature and prevalence of obesity in the Spanish population: The Di@bet.es study.](#) Valdés S, Maldonado-Araque C et al. *Obesity* (Silver Spring). 2014 Nov;22(11):2328-32. doi: 10.1002/oby.20866.

[Downregulation of G protein-coupled receptor kinase 2 levels enhances cardiac insulin sensitivity and switches on cardioprotective gene expression patterns.](#) Lucas E, Jurado-Pueyo M et al. *Biochim Biophys Acta*. 2014 Dec;1842(12 Pt A):2448-56. doi: 10.1016/j.bbadi.2014.09.004.